

Tectonic and eustatic controls on the distribution of sandstone and hot mudstone: the Goodrich-Shaftesbury transition, Upper Albian and Lower Cenomanian, NE British Columbia

Piotr J. Angiel*, University of Western Ontario, London, Canada

pangiel@uwo.ca

and

A. Guy Plint, University of Western Ontario, London, Canada

gplint@uwo.ca

GeoConvention 2012: Vision

Summary

During the Late Albian and Early Cenomanian, deposition of the Goodrich Sandstones and their mudstone equivalent (the Shaftesbury Formation) took place during a time of overall rapid, but variable subsidence rate and relative sea level rise. An allostratigraphic analysis of the proximal to distal facies variability showed that a high accommodation rate trapped shoreface and coastal plain deposits near the western margin of the basin. Only during times of slower subsidence or possible eustatic fall were sandstones were able to prograde up to ~70 km to the west, beyond which, the entire succession is mudstone-dominated. Only mudstones above the Base of Fish Scale Marker (BFSM) have elevated radioactivity, and this marker can be traced to a major transgressive event at the western margin of the basin. This transgression is interpreted to have led to the mixing of the cooler and less saline Boreal with the warmer and more saline Tethyan water-masses. Resultant density stratification appears to have caused bottom-water anoxia and hence preservation of abundant organic matter.

Introduction

Well-exposed sections in the B.C. Foothills, coupled with a large well-log database in the adjacent Deep Basin makes this proximal part of the foreland basin a natural laboratory in which to investigate the relationship between facies stacking patterns, stratal geometry, and possible tectonic and eustatic controls. At a basin scale, the detailed relationship between the lithostratigraphic units of northeastern British Columbia (the Hasler, Goodrich and Cruiser formations) and their equivalents in Alberta (the Shaftesbury Formation) have remained enigmatic since the pioneer work of Wickenden and Shaw (1943) and Stott (1968). Lithostratigraphy worked well for outcrop mapping in British Columbia, but on a regional scale, the lithostratigraphic divisions cannot be extended far to the east because of both facies diachroneity and lateral variability. Therefore, new allostratigraphic correlations, based on tracing allostratigraphic discontinuities (mostly flooding surfaces) is necessary. The allostratigraphic approach of Roca *et al.* (2008) illustrated how time markers cross different lithostratigraphic units from the Canada - USA border to the Alberta - British Columbia border. In this presentation, we now extend this correlation to the proximal foredeep in order to illustrate the lateral relationship between relatively sand-rich proximal facies (Hasler, Goodrich, Cruiser fms.) and more mud-rich distal facies (Shaftesbury Formation).

Theory and Method

The rocks in the studied interval are characterized by strongly cyclical, sandier-upward transgressive-regressive successions, typically 5-20 m thick. The flooding surfaces that bound these successions are readily traceable in subsurface, sometimes for > 400 km, and can be considered to approximate time lines. This proxy chronostratigraphy provides a basis for facies and paleogeographic mapping. Isopach maps constructed for allostratigraphic units (effectively time slice maps of subsidence) allowed reconstruction of the subsidence history in three dimensions, on small time-steps. Paleogeographic maps for each time slices makes it possible to trace, with high resolution, changes in facies distributions in response to subsidence and eustasy.

For this study gamma ray and/or resistivity logs from 435 wells were used to construct a grid of 20 cross sections with a spacing of ~20 km. These wells formed the basis for an allostratigraphic correlation grid over an area of about 50,000 km². Twenty one outcrop sections totaling 4090 m have been investigated. Effort was made to correlate well logs with nearby outcrop sections in order to establish the nature of facies and bounding surfaces that were defined and mapped in subsurface.

Results

Overall geometry

Allomembers show no clinoforms geometry, nor do they lap out onto a single discrete surface. Instead, each allomember fines and gradually thins eastward away from the orogen-proximal part of the foredeep. The sedimentological and stratigraphic evidence indicates that the proximal foredeep was a shallow, low-gradient, storm-dominated ramp. Sand was transported, primarily by combined flows, for up to 70 km offshore. The more distal part of the ramp accumulated mud. There is no evidence for the development of deep water (i.e. > ~30-40 m), and therefore the accommodation:supply ratio appears to have been near balance throughout deposition of the succession.

Sandstone – mudstone transition

Sandstone bodies comprising the Goodrich Formation were deposited in the most proximal foredeep, over about 1 My. Within the formation, eight major sandstone tongues, capped by flooding surfaces, can be recognized. Sandstones are stacked vertically in the west, where about 450 m of nearshore and coastal plain deposits are observed. An aggradational stacking pattern and a pronounced wedge-shaped geometry suggest that these units were deposited at a time of rapid subsidence and relative sea level rise. Lateral facies change is dramatic, and the progradational limit of inner-shelf sandstone (rippled heterolithic facies) never extends more than ~60-70 seaward of the paleoshoreline. It is interpreted that each progradational sandstone tongue was deposited during a phase of slower relative sea level rise, perhaps linked to periods of eustatic fall. It is remarkable that the Goodrich Sandstones vanish in about 70 km from the paleoshoreline, beyond which, the entire succession is represented by about 200 m of mudstone (the Shaftesbury Formation).

The Base of Fish Scales Marker and the origin of hot mudstones

The Base of Fish Scale Marker (BFSM) has a very prominent 'hot' signature in gamma ray logs, and is a basin-wide log marker. Towards the northwest, in British Columbia, the radioactive signature of the BFSM gradually fades. A chert-pebble veneered erosion surface underlies the BFSM and represents the most prominent sea level fall in the Goodrich time interval. The distribution of pebbles indicates that relative sea level fall caused the shoreline to move at least 130 km to the east. The overlying

transgressive mudstones have a very high radioactivity (>150 API). It is notable that hot mudstone is not present below the BFSM surface whereas above, there are two horizons with elevated radioactivity. In the upper horizon, the lateral transition from shelf sandstones into radioactive mudstone takes place over a distance of only about 50 km. It is interpreted that deposition of hot mudstone was linked to the major transgression that connected the Boreal and Tethyan Ocean and which caused bottom-water anoxia, due to density or salinity stratification (Schröder-Adams *et al.* 1996).

Conclusions

The Hasler, Goodrich and Cruiser formations are relatively sandstone-rich, basin-margin facies that were deposited at a time of rapid tectonic subsidence and relative sea level rise. The high accommodation rate in the western, proximal foredeep trapped coarser sediment and caused sandy, nearshore facies to aggrade vertically. A high resolution allostratigraphic approach permitted recognition of changes in stratal geometry. It appears that, at times of somewhat lower subsidence rate, shorelines were able to prograde further seaward, building more tabular-shaped rock bodies. Nevertheless, sand appears to have been transported only ~60-70 km seaward. An elevated radioactivity level is observed only above a pebble-veneered, formerly-subaerial unconformity that underlies the Base of Fish Scale Marker. This unconformity is blanketed by highly radioactive organic mudstones that were deposited following a major sea level rise that is interpreted to have allowed connection between the Boreal and Tethyan water masses, leading to density stratification, bottom-water anoxia, and preservation of organic matter.

This study provides clear confirmation of theoretical foreland basin models (e.g. Jordan and Flemings, 1991) that predict nearshore trapping of coarse clastics during times of rapid flexural subsidence, and the domination of the offshore portion of the basin by mud. Some of the offshore mudstone units are organic-rich, and have the potential to source hydrocarbons and host shale gas. Therefore, it is important to understand the tectonic and oceanographic conditions under which organic-rich mudstones may be deposited, and to understand the dispersal processes and facies variability within mud-rich facies across a shallow, low-gradient shelf.

Acknowledgements

Angiel acknowledges partial support through a Geoscience BC grant, from an SEPM Student Assistance Grant and from the AAPG Grants-In-Aid program. Plint acknowledges Discovery Grant support from the Natural Sciences and Engineering Research Council of Canada. We thank Divestco Ltd. for access to digital well-logs, and Imperial Oil Ltd. for donation of a microfiche log library. Capable field assistance was provided by Robin Buckley, Meriem Grifi, Magda Hrnkova, Tessa Plint, Joel Shank and Kristin Smith.

References

- Jordan, T.E., and Flemings, P.F., 1991, Large-scale stratigraphic architecture, eustatic variation, and unsteady tectonism: A theoretical evaluation. *Journal of Geophysical Research*, **96B4**, 6681-6699.
- Roca, X., Rylaarsdam, J.R., Zhang, H., Varban, B.L., Sisulak, C.F., Bastedo, K., and Plint, A.G., 2008, An allostratigraphic correlation of Lower Colorado Group (Albian) and equivalent strata in Alberta and British Columbia, and Cenomanian rocks of the Upper Colorado Group in southern Alberta: *Bulletin of Canadian Petroleum Geology*, **56**, 259-299.

- Schröder-Adams, C.J., Leckie, D.A., Bloch, J., Craig, J., McIntyre, D.J., and Adams, P.J., 1996, Paleoenvironmental changes in the Cretaceous (Albian to Turonian) Colorado Group of Western Canada: microfossil, sedimentological and geochemical evidence: *Cretaceous Research*, **17**, 311–365.
- Stott, D.F., 1968, Lower Cretaceous Bullhead and Fort St. John groups, between Smoky and Peace Rivers, Rocky Mountain Foothills, Alberta and British Columbia: *Geological Survey of Canada Bulletin*, **152**, 279 p.
- Wickenden, R.T.D., and Shaw, G., 1943, Stratigraphy and structure in Mount Hulcross-Commotion Creek map-area, British Columbia: *Geological Survey of Canada Preliminary Paper*, **43**, 1-13.